

Claims

1. A method for treating a human with a stimulant dependency, said method comprising administering to said human pramipexole in a dose ranging from 1.5 mg/day to 6.0 mg/day.
2. The method of claim 1, wherein said stimulant dependency involves a stimulant craving.
3. The method of claim 1, wherein said stimulant is cocaine.
4. The method of claim 1, wherein said pramipexole is administered in a dose ranging from 1.5 mg/day to 4.5 mg/day.
5. A method of treating a cocaine craving in a human, said method comprising administering to said human pramipexole in a dose ranging from 1.5 mg/day to 6.0 mg/day.
6. The method of claim 5, wherein said pramipexole is administered in a dose ranging from 1.5 mg/day to 4.5 mg/day.
7. A method for treating a human with a stimulant dependency, said method comprising administering to said human pramipexole, wherein said pramipexole is administered intranasally.
8. The method of claim 7, wherein said stimulant dependency involves a stimulant craving.
9. The method of claim 8, wherein said stimulant is cocaine.

10. The method of claim 7, wherein said pramipexole is administered in a dose ranging from 1.5 mg/day to 6.0 mg/day.

11. The method of claim 10, wherein said pramipexole is administered in a dose ranging from 1.5 mg/day to 4.5 mg/day.

12. A method of treating a cocaine craving in a human, said method comprising administering to said human pramipexole, wherein said pramipexole is administered intranasally.

13. The method of claim 12, wherein said pramipexole is administered in a dose ranging from 1.5 mg/day to 6.0 mg/day.

14. The method of claim 13, wherein said pramipexole is administered in a dose ranging from 1.5 mg/day to 4.5 mg/day.